Common name:African Green Wood Hoopoe	Scientific name: Phoeniculus Purpureus		Year first used: 2006	Diet last updated: 11/11/2013	Institution: Private Aviculturist. Gary Bralsford
Number of individuals diet fed to: 1 pair			Have birds bred whilst on this diet? Yes		
Diet Ingrediant	Amount per day (g)	Or standard m	easure	Preparation	
Black Cricket medium		12		-	
Brown Cricket large		12			
Morio worms			6		
Regular mealworms			egg cup full		
Waxworms			egg cup full		
Large Locust			4		
Waxmoths			hand full		
Mince Meat		de	esert spoon full		
Day old Chick			1	chopped into 1/2" pie	eces
Bird of Prey Diet			scoop		
Purina Dog Food (meat)		SCOO	p (desert spoon)	soaked & drained	
carrots & broccoli			<u> </u>	Diced	
C15 Pellets			scoop		
Berries (wild in season)			3333	picked fresh	
Nectar (kiezebrink sunbird)				mixed twice a day	
Honey Water				mixed twice a day	
Tioney Tracer				mixed times a day	
Diet supplements used every other feed every we		unt: Nutrabol	, Bee Plllen , Bird of F	I Prey Calcium Powd	er , fed each on
Feeding method: All food is given on feed trays. Insects fed in a deep bucket to prevent escape					
Seasonal changes, pre reduced. Reduce livefood	<u> </u>	-	Livefood is increased	in breeding seaso	n . Nectar is
Dietary enrichment: Wil	d Berries and Hone	y water gives	a boost after the bree	ding season , Meat	t is given more in the
winter months .					
Articles or references used to formulate this diet: Birds Of Africa . Taxon Advisory Group , and personal data from my experiences .					
Diet related health issues whilst using the above diet: Lack of animal protein can cause rickets in young from early nestings . Adults thrive from varied diet					